

REFERENCE

- Kathiresan K. & Duraisamy. (2005). Current issue of microbiology. *ENVIS Centre Newsletters*. vol. 4, pp. 3–5.
- Haefner B. (2003). Drugs from the deep: marine natural products as drug candidates. *Drug Discovery Today*. vol. 8, pp. 536–544.
- Yoshie Y, Wand W, Hsieh YP, Suzuki T. (2002). Compositional difference of phenolic compounds between two seaweeds, *Halimeda* spp. *J Tokyo univ Fish*. 88:21–24.
- Ponce NMA, Pujol CA, Damonte EB. (2003). Fucoidans from the brown seaweed *Adenocystis utricularis*: Extraction methods, antiviral activity and structural studies. *Carbohydr Res*. 338: 153–165.
- Riou D, Collic-Jouault S, Du Sel DP. (1996). Antitumor and antiproliferative effects of a fucan extracted from *Ascophyllum nodosum* against a non-small-cell bronchopulmonary carcinoma line. *Anticancer Res*. 16: 1213–1218.
- Jurd KM, Rogers DJ, Blunden G, McLellan DS. (1995). *J. Appl. Phycol*. 7 3339–3345.
- Sheng JC, Yu F, Xin ZH, Zhao LY, Zhu XJ, Hu QH. (2007). Preparation, identification and their antitumor activities in vitro of polysaccharides from *Chlorella pyrenoidosa*. *Food Chem*.105:533–9.
- Zhou GF, Sun YP, Xin H, Zhang Y, Li ZE, Xu ZH. (2004). In vivo antitumor and immunomodulation activities of different molecular weight lambda-carrageenans from *Chondrus ocellatus*. *Pharmacol Res*. 50:47–53.